

Explore by station

Explore by bacteria type

FIND A STATION:
Times Square-42nd St, Theater District (A, C, E, ...)

Where Certain Types of Bacteria Were Found


Select a category below to see the stations where researchers found the associated types of bacteria.

- Antibiotic resistance
- Diarrhea
- Dysentery
- Food poisoning
- Heart-valve infections
- Italian cheese
- Kimchi and sauerkraut**

Leuconostoc citreum

Associated with **Kimchi and sauerkraut**
A species called *Leuconostoc citreum* is widely used in the fermentation of the Korean dish kimchi and is also used to ferment cabbage to make sauerkraut.

- Medical-device infections
- Meningitis
- Mozzarella cheese
- Oil cleanup
- Radiation resistance
- Respiratory ailments
- Sepsis
- Staph infections
- Sunscreen
- Swiss cheese
- Tetanus
- Toxic cleanup
- Urinary-tract infections



Share this view:

Interactive by Martin Burch, Chris Canipe, Madeline Farbman, Rachel Feirman and Robert Lee Hotz

Notes: Two stations were closed in Brooklyn and could not be sampled: Central Av (served by M trains) and Kings Highway (B and Q trains). In some cases, data from different subway lines have been combined at station complexes.

Sources: PathoMap Project, Weill Cornell Medical College, American Society for Microbiology, U.S. Centers for Disease Control and Prevention, National Institutes of Health, U.S. National Library of Medicine, MSDSonline, MicrobeWiki, Metropolitan Transportation Authority, Mapbox, OpenStreetMap, Ontodia